

The Plasma Structure of the Southwestern Region of the Cygnus Loop with the XMM-Newton Observatory

Hiroyuki Uchida¹, Hiroshi Tsunemi¹, Satoru Katsuda^{1 2} and Masashi Kimura¹

uchida@ess.sci.osaka-u.ac.jp, tsunemi@ess.sci.osaka-u.ac.jp,
katsuda@ess.sci.osaka-u.ac.jp, mkimura@ess.sci.osaka-u.ac.jp

ABSTRACT

We observed the southwestern region of the Cygnus Loop in two pointings with *XMM-Newton*. The region observed is called the “blow-out” region that is extended further in the south. The origin of the “blow-out” is not well understood while it is suggested that there is another supernova remnant here in radio observation. To investigate the detail structure of this region in X-ray, we divided our fields of view into 33 box regions. The spectra are well fitted by a two-component nonequilibrium ionization model. The emission measure distributions of heavy elements decrease from the inner region to the outer region of the Loop. Then, we also divided our fields of view into 26 annular sectors to examine the radial plasma structure. Judging from metal abundances obtained, it is consistent with that the X-ray emission is the Cygnus Loop origin and we concluded that high- kT_e component (~ 0.4 keV) originates from the ejecta while low- kT_e component (~ 0.2 keV) is derived from the swept-up interstellar medium. The flux of low- kT_e component is much less than that of high- kT_e component, suggesting the ISM component is very thin. Also, the relative abundances in the ejecta component shows similar values to those obtained from previous observations of the Cygnus Loop. We find no evidence in X-ray that the nature of the “blow-out” region originated from the extra supernova remnant. From the ejecta component, we calculated the masses for various metals and estimated the origin of the Cygnus Loop as the core-collapse explosion rather than the Type Ia supernova.

Subject headings: ISM: abundances — ISM: individual (Cygnus Loop) — supernova remnants — X-rays: ISM

¹Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

²NASA Goddard Space Flight Center, Code 662, Greenbelt, MD 20771

1. Introduction

The Cygnus Loop is one of the brightest Supernova Remnant (SNR) in the X-ray sky. Its age is estimated to be $\sim 10,000$ yrs (Blair et al. 2005). Since the distance is comparatively close to us (540 pc; Blair et al. 2005), the apparent size is quite large ($2.5^\circ \times 3.5^\circ$; Levenson et al. 1997), which enables us to study the plasma structure of the Loop in detail.

Although the Cygnus Loop is an evolved SNR, a hot plasma is still confined inside the Loop (Hatsukade & Tsunemi 1990). Miyata et al. (1998) observed the Loop with the *Advanced Satellite for Cosmology and Astrophysics* (ASCA), and detected the strong highly-ionized Si-K, S-K line and Fe-L line near the center of the Cygnus Loop. They concluded that the hot plasma, a “fossil” of the supernova explosion, left in the core of the Loop. Tsunemi et al. (2007) (hereafter TKNM07) observed the Cygnus Loop along the diameter from the northeast (NE) to the southwest (SW) with *XMM-Newton* and studied the radial plasma structure. From the spectral analysis, they showed that the Cygnus Loop consists of two component plasma. They concluded that the low- kT_e component originating from the interstellar medium (ISM) surrounds the high- kT_e component originating from the ejecta. In addition, they measured the metal abundances of the high- kT_e component and showed the metal distribution of the ejecta. The results indicate that the abundances are relatively high (~ 5 times solar) and each element is nonuniformly distributed: Si, S and Fe are concentrated in the inner region while the other elements such as O, Ne and Mg are abundant in the outer region. They also estimated the progenitor star’s mass to be $15 M_\odot$.

The Cygnus Loop is a typical shell-like SNR; this structure is thought to be generated by the cavity explosion (Levenson et al. 1997). The Cygnus Loop is almost circular in shape, however, we can see some breakout in the SW. It is called the “blow-out” region (Aschenbach & Leahy 1999). The origin of the “blow-out” is not well understood. Aschenbach & Leahy (1999) have explained this extended structure as a breakout into a lower density ISM. On the other hand, Uyaniker et al. (2002) suggested the existence of a secondary SNR (named G72.9-9.0) in the south from a radio observation and some other radio observations support this conclusion (Uyaniker et al. 2004; Sun et al. 2006).

Our observations were performed in a direction from the Cygnus Loop center toward the south “blow-out” region. In this paper, we report the result of the spectral analysis and discuss about the plasma structure of this region.

2. Observations

We performed two pointing observations of the SW region of the Cygnus Loop with the *XMM-Newton* observatory. TKNM07 have observed from the NE to the SW along the diameter. Then, we intended to expand our observation southward from the center. Figure 1 left panel shows the X-ray surface brightness map of the Cygnus Loop obtained with *ROSAT* High Resolution Imager (HRI) on which we showed our Field of View (FOV) of the EPIC MOS by solid white circles. We call the north observation for Position-8 (Pos-8) and the south observation for Position-9 (Pos-9). If there exists a secondary SNR in SW as Uyaniker et al. mentioned, our whole FOV overlaps the SNR whose center is roughly located at the south in Pos-9. Figure 1 right shows a three-color X-ray image of our FOV using *XMM-Newton* EPIC MOS 1 and 2 data after correcting for exposure and vignetting effects. Red, green and blue correspond to the energy ranges of 0.3-0.5 keV, 0.5-0.7 keV and 0.7-3.0 keV, respectively. Figure 2 shows the MOS broad-band image for the 0.3-3 keV range. The white X shows the center of the G72.9-9.0 estimated by Uyaniker et al. (2002).

Both observations were performed on May 13 2006, during the *XMM-Newton* AO-5 observing cycle. The total exposure time was both ~ 10 ks. In order to exclude the background flare events, we considered the time intervals when the count rates were high as flare events, and eliminated them from our analysis. The pn data were almost unusable due to the flare. Therefore we only used the data obtained with the EPIC MOS for our analysis. The effective exposure times for each observation were 6.5 ks (Pos-8) and 3.6 ks (Pos-9), respectively. Both data were taken using the medium filters and the prime full-window mode. All the data were processed with version 7.1.0 of the *XMM* Science Analysis System (SAS). For the background subtraction, we employed a blank-sky observations prepared by Read & Ponman (2003). The point-like sources were excluded using the SAS task edetect chain for the spectral analysis. As a result, one and four point-like sources are detected in Pos-8 and Pos-9, respectively. Two of the point sources in Pos-9 were observed by Miyata et al. (2001) and named AX J2049.6+2939 and AX J2050.0+2914, respectively.

3. Spectral Analysis

3.1. Two-component VNEI Model

Figure 3 shows the spectra for Pos-8 and 9 summed over the entire FOV. We can see some emission lines such as O He α , O Ly α , the Fe L complex, Ne He α , Mg He α , and Si He α , while S line is not seen here.

Firstly, we fitted each spectrum by single-component non-equilibrium ionization (VNEI) model. We employed **wabs** (Morrison & McCammon 1983) and **vnei** (NEI ver.2.0; Borkowski et al. 2001) in XSPEC version 12.3.1 (Arnaud 1996). In the model, the abundances of O, Ne, Mg, Si, and Fe were free while we set the abundances of C and N equal to O, S equal to Si, Ni equal to Fe, and other elements fixed to their solar values (Anders & Grevesse 1989). Other parameters were all free such as the electron temperature kT_e , the ionization timescale τ (a product of the electron density and the elapsed time after the shock heating), and the emission measure ($EM = \int n_e n_H dl$, where n_e and n_H are the number densities of hydrogen and electrons and dl is the plasma depth). We also set the column density N_H free. From the best-fit parameters, we found that the value of kT_e (~ 0.4 keV) is higher than that of the result at the NE rim obtained from *Suzaku* observations (~ 0.2 keV) (Katsuda et al. 2008a). Also the metal abundances such as Si (~ 1.0) and Fe (~ 0.4) show about two times higher values than those of the NE rim. These facts suggest that the X-ray emission in Pos-8 and 9 mainly consists of the high- kT_e component which was explained in section 1. However, the values of reduced χ^2 are 6.9 and 3.6 in Pos-8 and 9, respectively. The model is not enough to fit the data due to the simplicity of the model. Therefore, we intended to add the extra component to the VNEI model.

From the standpoint of the SNRs evolution, the X-ray emission from the SNRs have the two different origin. The blast wave from the supernova explosion sweeps the ambient medium, while the reverse shock propagates into the ejecta. Each shock wave heats up the swept-up ISM and the ejecta respectively. The shock-ISM interaction also produces the reflected shock which moves back through previously swept up ISM (Hester et al. 1994). Because of this, the X-ray spectra of the evolved SNR such as the Cygnus Loop should have a complicated structure. From the earlier observation of the NE to the SW regions, TKNM07 proposed the plasma structure of the Cygnus Loop as follows: the high- kT_e ejecta component is surrounded by the low- kT_e ISM component. They found that the spectra from most regions of the Cygnus Loop consist of the two-component VNEI model. Thus we also employed two-component VNEI model which has two different electron temperatures. We found that this model cannot reach the physically meaningful results by setting all the parameters free. Therefore, in the low- kT_e component, we fixed the metal abundances to the values from the NE rim observations (Uchida et al. 2006). Other parameters were set free such as kT_e , τ , and EM. In the high- kT_e component, we set all parameters to those of the single- kT_e VNEI model as explained in the above paragraph. As a result, the values of the reduced χ^2 remain almost unchanged: 6.9 to 6.5 and 3.6 to 3.4 in Pos-8 and 9, respectively. These large values are due to the fact that we took the spectra summed over the entire FOV in which there is a lot of structure. Therefore we divided our FOV into several regions for the spectral analysis. Although we employed the constant temperature, plane-parallel shock

plasma model, **vpshock** instead of **vnei**, the best-fit parameters were almost unchanged and the values of the reduced χ^2 were not significantly improved.

3.2. Spatially Resolved Spectral Analysis

From Fig. 2, we can see a lot of structures within Pos-8 and 9. For example, there is a region of high surface brightness at each center of Pos-8 and 9 even after correcting the vignetting effect. We notice that they are different in color in Fig. 1 right, which shows that the plasma temperatures are different from each region. In order to investigate the detail plasma structure, we divided our FOV into a number of box regions for the spectral analysis. To equalize the statistics, we determined the box sizes such that each region has 7,500-15,000 photons for MOS 1 and 2. In this way, we have 33 regions (22 and 11 regions in Pos-8 and Pos-9, respectively). Figure 2 left panel shows the *XMM-Newton* MOS broad band image for the 0.3-3.0 keV range and box regions are shown in white lines.

To examine the plasma structure of Pos-8 and 9, we fitted 33 spectra extracted from box regions by the single- kT_e VNEI model and two- kT_e VNEI model, respectively. In the two- kT_e VNEI model, we fixed the metal abundances of the low- kT_e component to the result from the observations of the NE rim as explained in section 3.1. As in the case of the fit for each whole region, the values of the reduced χ^2 are improved ~ 1.6 to ~ 1.3 and ~ 1.4 to ~ 1.1 in Pos-8 and 9, respectively. The F-test probability ($> 99\%$) shows that it is reasonable to add the extra low- kT_e VNEI model in more than half of the regions. The best-fit parameters are shown in Figure 4 as the maps of the best-fit parameters. The averaged temperature of high- and low- kT_e component are ~ 0.4 keV and ~ 0.2 keV, respectively. However, the low- kT_e temperatures are determined only as the upper limit in several regions where the contribution of the low- kT_e component is quite low as shown in the EM_L map in Fig. 4.

We compared these parameters and EMs of heavy elements with the results of Katsuda et al. (2008b). They observed the Cygnus Loop in seven pointings from the NE to the SW with *Suzaku* and showed the best-fit parameters using the two- kT_e VNEI model. One of their observation regions (named P16) is next to the NE part of Pos-8 (see Fig. 1). From the results of P16 observations, the temperature of the high- and low- kT_e component are 0.4 keV and 0.2 keV. These values are similar to our results. Katsuda et al. concluded that the emission of the high- kT_e component comes from the ejecta of the Cygnus Loop. Then we compared the EMs of O [=C=N], Ne, Mg, Si [=S], and Fe [=Ni] between in P16 and our FOV as shown in Table 3. Katsuda et al. (2008b) showed that each EM in their FOV reduces from the center to the outer region of the Loop. From Table 3, we found that this trend is also seen from P16 to our FOV.

Then, we determined the spectral extraction regions in different way to investigate the plasma structure from the inner side of the Loop to the outside.

We divided our FOV into two paths: the east path and west path. Then we divided several annular sectors as shown in Fig. 2 right. To compare our analysis with that of TKNM07, we set the annular center on $20^h51^m34.7^s, 31^\circ00'00''$ (J2000). In order to equalize the statistics, we determined the annular widths such that each sector has at least $\sim 10,000 \pm 1,000$ photons. In this way, we have 26 annular sectors (16 and 10 sectors in the east and the west path, respectively) whose angular distances from the center are from 35 arcmin to 95 arcmin. The width ranges from 1 arcmin to 6.5 arcmin.

Figure 5 shows the example of the spectrum from the sector at $R = 42.5'$, where R represents the angular distance from the center. The left and right panels show the best-fit curves with the single- and the two- kT_e VNEI model, respectively. The fitting parameters are set as explained in section 3.1. Dotted lines in Fig. 5 represent the individual model. From Fig. 5 right, we found that the contribution of the low- kT_e component is lower than that of the high- kT_e component. The best-fit parameters are shown in Table 1 and 2, respectively. The F-test probability ($> 99\%$) shows that it is reasonable to add the extra low- kT_e VNEI model in this sector.

Then, we analysed all other sectors in the same way. Figure 6 shows the radial plot of the values of χ^2 along the east path (top) and the west path (bottom). The single- kT_e VNEI model is shown in black, while the two- kT_e VNEI model is shown in red. From the results, we calculated the F-test probability and determined whether or not the extra component is needed for each sector. Applying the significance level of 99%, the extra component is not required at $47.5' < R < 75.0'$, and $77.5' < R < 95.0'$ along the east path, and $36.0' < R < 43.0'$, $47.0' < R < 65.0'$, and $85.0' < R < 95.0'$ along the west path. In other words, $\sim 60\%$ of our FOV requires the two- kT_e VNEI model. We find that the fit shown in Fig.5 (at $R = 42.5'$) is improved the most by using the two- kT_e VNEI model. Even in this sector, the contribution of the additional low- kT_e component is not so large.

4. Discussion

The first two panels of Fig. 4 show the temperature distributions of the two components based on the analysis in Fig. 2 left. Figure 7 shows the temperature distributions along the east path (top) and the west path (bottom) based on the analysis in Fig. 2 right. Black and red represent the low- kT_e temperature and the high- kT_e temperature. From Fig. 4 and 7, the averaged values of low- and high- kT_e temperature are ~ 0.2 keV and ~ 0.4 keV, respectively.

In this way, we clearly separated the high- kT_e component and the low- kT_e component just as the observation obtained in Katsuda et al. (2008b) and TKNM07. TKNM07 showed the temperature of the low- kT_e component is almost constant (~ 0.2 keV) along the diameter, while that of the high- kT_e component is different in NE (~ 0.6 keV) and SW (~ 0.4 keV). Since our FOV is very close to SW, our result shows smooth extrapolation from that of TKNM07 in SW rather than that in NE.

The third and fourth panels of Fig. 4 shows the EM distributions for each component. Although there are some structures seen in the EM of the high- kT_e component (EM_H) map, it is clear that the EM_H is higher in Pos-8 than that in Pos-9. The EM of the low kT_e component (EM_L) in all of our FOV are lower than those of EM_H . Figure 8 shows the radial profile of the EMs for each component. In this figure, we calculated the EMs as a function of R into 10 arcmin bin. The EM_L stays almost constant while it peaks around $R = 80'$. From the morphological point of view, the Cygnus Loop has an almost circular shape with a radius $\sim 80'$ except for the south “blow-out”. Then, it is suggested that the EM_L distribution reflects the rim brightening structure around the “blow-out” region. On the other hand, the value of EM_H gradually decreases from the center to the outer region. This decrease can be easily explained by assuming that the emission comes from the ejecta component filling inside the Cygnus Loop. Then we also measured the EMs of various heavy elements in the high- kT_e component such as O [=C=N], Ne, Mg, Si [=S], and Fe [=Ni] and compared them with the result of TKNM07. Figure 9 shows the EM distribution for these elements as a function of R . We also plot the results of Pos-2 to 6 (TKNM07) in the same panels. Although some structures are remaining in the annular regions as seen in Fig. 2, the radial distribution of each EM clearly shows the smooth extrapolation of TKNM07’s result. They showed the decrease of each EM from the center to the outer region and concluded that the high- kT_e component is derived from the Cygnus Loop ejecta. It is reasonable to understand that the EMs in Pos-8 and 9 shows a smooth connection to those in their FOV. Therefore we concluded that the high- kT_e component originates from the ejecta of the Cygnus Loop.

From the fitting parameters of the high- kT_e component, we calculated the abundances of ejecta component for various elements. Figure 10 shows the abundance ratios of heavy elements (Ne: black, Mg: red, Si [=S]: green, Fe [=Ni]: red) relative to O. From Fig. 10, we found that Si/O (~ 20) and Fe/O (~ 10) are heavily over abundant and Ne/O is ~ 2 , while Mg/O (< 1) is depleted. This tendency is kept throughout our observing region. The other observations of the ejecta in the Loop such as TKNM07 and Katsuda et al. (2008b) showed the similar results.

Uyaniker et al. (2002) and Sun et al. (2006) reported that the Cygnus Loop consists of two SNRs interacting with each other in the SW. Their main arguments are the

difference of the radio morphology and the polarization intensity between the main part of the Cygnus Loop and the south “blow-out” region. However, based on the X-ray data, we found that there is no evidence of the extra SNR within our FOV. If these SNRs are at the same distance as claimed by Uyaniker et al. (2002) and Sun et al. (2006), the smaller radius of the extra SNR, ~ 7 pc ($R/\sim 0.7^\circ$)($d/540$ pc), than that of the Cygnus Loop, ~ 13 pc ($R/\sim 1.4^\circ$)($d/540$ pc), strongly suggests that the extra SNR is younger than the Cygnus Loop is. If we employ the Sedov-Taylor solution, the temperature T of the extra SNR is

$$T \sim 1.8 \text{ keV} \left(\frac{E_0}{10^{51} \text{ ergs}} \right) \left(\frac{n}{1 \text{ cm}^{-3}} \right)^{-1} \left(\frac{R}{7 \text{ pc}} \right)^{-3} \quad (1)$$

where E_0 and n are the explosion energy and the surrounding medium density of the extra SNR, respectively. Therefore, the temperature of the extra SNR should be significantly higher than that of the Cygnus Loop. However, we found no sign of such high temperature plasma. The spectra from all regions are almost represented by a single- kT_e VNEI model (~ 0.4 keV). If we add an extra component, we found in section 3 that the extra component shows low temperature rather than high temperature. Furthermore, from Fig. 7, the temperatures of each component are in good agreement with those obtained in other regions of the Loop (TKNM07; Katsuda et al. 2008b). This result suggests that the X-ray emission from the SW region mainly comes from the Cygnus Loop. If the secondary SNR exists in the SW at the same distance, the contribution of the X-ray emission to the spectra is much less than that of the Cygnus Loop. We cannot rule out the possibility that the extra SNR exists far side of the Cygnus Loop. However, even if that is the case, the fact remains that the spectrum from our FOV mainly consists of the ejecta and the ISM components of the Cygnus Loop. As a result, we find no evidence in X-ray that there exists the second SNR at the same distance to the Cygnus Loop.

Then, we can estimate the mass of the progenitor star of the Cygnus Loop from the EMs of the high- kT_e component, assuming that all these emissions come from the Loop. Then, we multiplied the EMs by the area of each annular sector and integrated the EMs along the path. In this way, we obtained the emission integral ($EI = \int n_e n_X dV$, where dV is the X-ray-emitting volume) for O, Ne, Mg, Si, and Fe. Table 4 shows the calculated EI of each element. To compare our data with the supernova explosion models, we calculated the ratios of Ne, Mg, Si and Fe relative to O. Figure 11 shows the number ratios of Ne, Mg, Si and Fe relative to O of the ejecta component. We also plotted the result from TKNM07, the core-collapse models (Woosley & Weaver 1995) for various progenitor masses and Type Ia supernova models (Iwamoto et al. 1999) for comparison. The type Ia supernova yields more Fe than our results but less Ne. On the contrary, the number ratios of Si and Fe are higher than those of any core-collapse models, which attributes to the fact that our calculations

are derived from the small part of the Cygnus Loop. These models were calculated under the assumption of the symmetric explosion. However, TKNM07 and Katsuda et al. (2008b) reported the asymmetry of the metal distribution of the ejecta component: Si and Fe were more abundant, while Ne and Mg were less abundant in the SW rather than that in the NE. These results support an asymmetric explosion of the progenitor star. Our FOV is close to the SW region in TKNM07 rather than the NE region. Taking into account the effect of the asymmetric structure, we support the idea that the Cygnus Loop originates from the core-collapse explosion rather than the Type Ia supernova.

5. Conclusion

We observed the SW region of the Cygnus Loop with *XMM-Newton*. To examine the plasma structure, we divided our FOV in two different ways: 33 box sectors and 26 annular sectors. We fitted the spectrum extracted from each region with two- kT_e VNEI model. The plasma structure of the low- kT_e component and that of the high- kT_e component are quite different from each other: each temperature is ~ 0.2 keV and ~ 0.4 keV for the former and the latter, respectively.

The EM distribution of the low- kT_e component suggest the rim brightening structure, while that of the high- kT_e component monotonously decreases from the center of the Loop to the outside. In the high- kT_e component, the abundances of Si and Fe are relatively high compared to those of Ne and Mg. The distributions of EMs as well as the relative abundances in the high- kT_e component match the view that the low- and high- kT_e components, respectively, originate from the ISM and the ejecta of the Cygnus Loop, which was derived by earlier observations such as TKNM07 or Katsuda et al. (2008b). We found that the emission from this ISM component is relatively weak. This suggests that the thickness of the shell is thin in Pos-8 and 9. We also calculated the relative abundances of Ne, Mg, Si, and Fe to O in the ejecta component for the entire FOV, and estimated the origin of the Cygnus Loop as the core-collapse explosion rather than the Type Ia supernova. We found no evidence in X-ray that the nature of the “blow-out” region originated from the extra SNR.

We thank H. Kosugi’s careful reading of the manuscript. This work is partly supported by a Grant-in-Aid for Scientific Research by the Ministry of Education, Culture, Sports, Science and Technology (16002004). This study is also carried out as part of the 21st Century COE Program, ‘*Towards a new basic science: depth and synthesis*’. H.U. and S.K. are supported by JSPS Research Fellowship for Young Scientists.




Fig. 1.— *Left*: X-ray surface brightness map of the Cygnus Loop obtained with the *ROSAT* High Resolution Imager (HRI). The dotted circles and solid circles represent the FOV of TKNM07 and our observations (Position-8 and 9), respectively. The dotted square represents the FOV of *Suzaku* observation named P16 (Katsuda et al. 2008b). *Right*: Three-color X-ray image for Pos-8 and 9 using EPIC MOS 1 and 2 data.




Fig. 2.— *Left*: *XMM-Newton* MOS broad-band image for the 0.3-3 keV range. White lines represents the spectral extraction regions. The green circles show the point-like source regions excluded from our spectral analysis. The white X shows the center of the G72.9-9.0 estimated by Uyaniker et al. (2002) *Right*: Same as the left panel, but for the different spectral extraction regions.




Fig. 3.— *Left*, MOS 1 (black) and MOS 2 (red) spectrum for Pos-8 which are summed over the entire FOV. The best-fit curves are shown as solid lines. The dotted lines show individual component of the model. The lower panel shows the residual. *Right*, same as the left panel, but for Pos-9.




Fig. 4.— Maps of the best-fit parameters. EM_H and EM_L mean the emission measure of the high- and low- kT_e component, respectively. Last five panels show the EMs of O [=C=N], Ne, Mg, Si [=S], and Fe [=Ni] for the high- kT_e component in units of 10^{14}cm^{-5} . The values of kT_e and EM_[H,L] are in units of keV and 10^{18}cm^{-5} , respectively.




Fig. 5.— Example spectra at $R = 42.5'$. The solid line of each panel shows the best-fit curve with the single- kT_e VNEI model and the two- kT_e VNEI model, respectively. Each lower panel shows the residual. The dotted lines of the right panel show individual component of the two- kT_e VNEI model.




Fig. 6.— Radial plot of the values of χ^2 as a function of R along the east path (top) and the west path (bottom). The single- and two- kT_e VNEI model are shown in black and red, respectively. The degrees of freedom are all ~ 130 .




Fig. 7.— Temperature distributions of the two components as a function of R along the east path (top) and the west path (bottom). Red shows the high- kT_e component, while black shows the low- kT_e component.




Fig. 8.— *Left*: EM_L distribution as a function of R . *Right*: Same as the left, but for EM_H .




Fig. 9.— EM distributions for various metals (O [=C=N], Ne, Mg, Si [=S], and Fe [=Ni]) in the ejecta. Black and red show the west path and east path, respectively. Green shows the result of TKNM07 taken from Pos-2 to Pos-6.




Fig. 10.— Distributions of relative abundance of heavy elements to O are shown as a function of R . Si/O, Fe/O, Ne/O, and Mg/O are shown in green, blue, black, and red, respectively. The results of east path and those of west path are plotted in the same color.

Table 1: Spectral Fit Parameters

Parameter	Pos-8
N_H [10^{20}cm^{-2}]	4 ± 1
kT_e [keV]	0.46 ± 0.02
O(=C=N)	0.04 ± 0.01
Ne	0.09 ± 0.01
Mg	0.04 ± 0.02
Si	0.79 ± 0.16
Fe(=Ni)	0.33 ± 0.03
$\log \tau$	10.58
EM ^a [10^{18}cm^{-5}]	2.54 ± 0.56
χ^2/dof	187/133

Note. — Other elements are fixed to solar values. The errors are in the range $\Delta\chi^2 < 2.7$ for one parameter.

^aEmission Measure, $\int n_e n_H dl$




Fig. 11.— Number ratios of Ne, Mg, Si, and Fe relative to O of the high- kT_e component, estimated for the entire Loop (*solid line*). Dotted and solid black lines show the result from TKNM07's FOV and that from our FOV. Dotted red lines represent the CDD1 and W7 Type Ia supernova models of Iwamoto et al. (1999). Dotted blue, light blue, magenta, and green lines represent core-collapse models with progenitor masses of 12, 13, 15, 20 $M_{\odot}$, respectively (Woosley & Weaver 1995).

Table 2: Spectral Fit Parameters

Parameter	Value
N_H [10^{20}cm^{-2}]	< 4
Low- kT_e component:	
kT_e [keV]	0.26 ± 0.07
C	0.27 (fixed)
N	0.10 (fixed)
O	0.11 (fixed)
Ne	0.21 (fixed)
Mg	0.17 (fixed)
Si	0.34 (fixed)
S	0.17 (fixed)
Fe(=Ni)	0.20 (fixed)
$\log \tau$	10.62
EM^a [10^{18}cm^{-5}]	< 0.57
High- kT_e component:	
kT_e [keV]	0.47 ± 0.02
O(=C=N)	0.06 ± 0.02
Ne	0.08 ± 0.03
Mg	0.06 ± 0.03
Si	0.65 ± 0.14
Fe(=Ni)	0.42 ± 0.05
$\log \tau$	11.29 ± 0.03
EM^a [10^{18}cm^{-5}]	1.88 ± 0.38
χ^2/dof	167/130

Note. — Other elements are fixed to solar values. The errors are in the range $\Delta\chi^2 < 2.7$ for one parameter.
^aEmission Measure, $\int n_e n_H dl$

Table 3: Comparison between the averaged EM for each element in our FOV and P16 (Katsuda et al. 2008b)

EM of Each Element			
(10^{14} cm^{-5})	P16	Pos-8	Pos-9
O(=C=N)	4.2	1.2	0.28
Ne	0.42	0.25	0.09
Mg	0.04	0.02	0.01
Si	1.7	0.54	0.22
Fe(=Ni)	1.1	0.41	0.12

Table 4: Calculated Emission Integrals ($= \int n_e n_X dV$) of the Cygnus Loop Ejecta

EI	
Element	(10^{52} cm^{-3})
O	1.34 ± 0.20
Ne	0.34 ± 0.03
Mg	0.02 ± 0.01
Si	1.02 ± 0.10
Fe	0.65 ± 0.03

REFERENCES

- Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Arnaud, K. A. 1996, in ASP Conf. Ser. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes (San Francisco: ASP), 17
- Aschenbach, B., & Leahy, D. A. 1999, *A&A*, 341, 602
- Blair, W. P., Sankrit, R., & Raymond, J. C. 2005, *AJ*, 129, 2268
- Borkowski, K. J., Lyerly, W. J., & Reynolds, S. P. 2001, *ApJ*, 548, 820
- Hatsukade, I. & Tsunemi, H. 1990, *ApJ*, 362, 566
- Hester, J. J., Raymond, J. C., & Blair, W. P. 1994, *ApJ*, 420, 721
- Iwamoto, K., Brachwitz, F., Nomoto, K., Kishimoto, N., Umeda, H., Hix, W. R., & Thielemann, F.-K. 1999, *ApJS*, 125, 439
- Katsuda, S., Tsunemi, H., Uchida, H., Miyata, E., Nemes, N., Miller, E. D., Mori, K., & Hughes, J. P. 2008, *PASJ*, 60, S115
- Katsuda, S., Tsunemi, H., Miyata, E., Mori, K., Namiki, M., Nemes, N. & Miller, E. D. 2008, *PSAJ*, 60, S107
- Levenson, N. A., Graham, J. R., & Walters, J. L. 1997, *ApJ*, 484, 304
- Minkowski, R. 1958, *Rev. Mod. Phys.*, 30, 1048
- Miyata, E., Katsuda, S., Tsunemi, H., Hughes, J. P., Kokubun, M., & Porter, F. S. 2006, *PASJ*, 59, S163
- Miyata, E., Ohta, K., Torii, K., Takeshima, T., Tsunemi, H., Hasegawa, T., & Hashimoto, Yasuhiro. 2001, *ApJ*, 550, 1023
- Miyata, E. & Tsunemi, H. 1999, *ApJ*, 525, 305
- Morrison, R. & McCammon, D. 1983, *ApJ*, 270, 119
- Sun X. H., Reich, W., Han, J. L., Reich, P., & Wielebinski, R. 2006, *A&A*, 447, 937
- Tsunemi, H., Katsuda, S., Norbert, N., & Miller, E. D. 2007, *ApJ*, 671, 1717
- Uchida, H., Katsuda, S., Miyata, E., Tsunemi, H., Hughes, J. P., Kokubun, M., & Porter, F. S. 2006, a poster presentation at Suzaku Conference (2006) held in Kyoto

Uyaniker, B., Reich, W., Yar, A., Kothes, R., & Fürst, E. 2002, *A&A*, 389, L61

Uyaniker, B., Reich, W., Yar, A., & Fürst, E. 2004, *A&A*, 426, 909

Woosley, S. E., & Weaver, T. A. 1995, *ApJS*, 101, 181